

Lecture 4

Lecture 4

- ▶ Sinusoids in continuous time: example

Lecture 4

- ▶ Sinusoids in continuous time: example
- ▶ Addition of sinusoids using phasors

Lecture 4

- ▶ Sinusoids in continuous time: example
- ▶ Addition of sinusoids using phasors
- ▶ Introduction to discrete-time sinusoids

Example: Continuous-Time Sinusoid

Example: Continuous-Time Sinusoid

$$x(t) = A \cos(\Omega t + \phi)$$

Example: Continuous-Time Sinusoid

$x(t) = A \cos(\Omega t + \phi)$ is such that

Example: Continuous-Time Sinusoid

$x(t) = A \cos(\Omega t + \phi)$ is such that

- ▶ $x(t) \geq 3.0$ for $(1/6)^{\text{th}}$ of each period

Example: Continuous-Time Sinusoid

$x(t) = A \cos(\Omega t + \phi)$ is such that

- ▶ $x(t) \geq 3.0$ for $(1/6)^{\text{th}}$ of each period
- ▶ it takes 0.02 seconds for the value of the sinusoid to drop from 3.0 to 0.0

Example: Continuous-Time Sinusoid

$x(t) = A \cos(\Omega t + \phi)$ is such that

- ▶ $x(t) \geq 3.0$ for $(1/6)^{\text{th}}$ of each period
- ▶ it takes 0.02 seconds for the value of the sinusoid to drop from 3.0 to 0.0
- ▶ $x(0) = -\sqrt{3}$ and the first derivative $x'(0)$ is positive

Example: Continuous-Time Sinusoid

$x(t) = A \cos(\Omega t + \phi)$ is such that

- ▶ $x(t) \geq 3.0$ for $(1/6)^{\text{th}}$ of each period
- ▶ it takes 0.02 seconds for the value of the sinusoid to drop from 3.0 to 0.0
- ▶ $x(0) = -\sqrt{3}$ and the first derivative $x'(0)$ is positive

Based on given information, determine A , Ω and ϕ .