

Program to calculate symmetric weights for storage of 4-vectors; array origin set to 1

We wish equilibrium points of $Cdv/dt = Wy - Gv + I$, $y=y(v)$

$$C1 := 5 \quad C2 := 5 \quad C3 := 5 \quad C4 := 5$$

$$G1 := 2 \quad G2 := 2 \quad G3 := 2 \quad G4 := 2$$

$$C := \begin{bmatrix} C1 & 0 & 0 & 0 \\ 0 & C2 & 0 & 0 \\ 0 & 0 & C3 & 0 \\ 0 & 0 & 0 & C4 \end{bmatrix} \quad C = \begin{bmatrix} 5 & 0 & 0 & 0 \\ 0 & 5 & 0 & 0 \\ 0 & 0 & 5 & 0 \\ 0 & 0 & 0 & 5 \end{bmatrix}$$

$$G := \begin{bmatrix} G1 & 0 & 0 & 0 \\ 0 & G2 & 0 & 0 \\ 0 & 0 & G3 & 0 \\ 0 & 0 & 0 & G4 \end{bmatrix} \quad G = \begin{bmatrix} 2 & 0 & 0 & 0 \\ 0 & 2 & 0 & 0 \\ 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 2 \end{bmatrix}$$

Initial choice of equilibrium points, to be later augmented to obtain symmetry

$$v1 := \begin{bmatrix} 0.5 \\ 0.25 \\ 0 \\ 0 \end{bmatrix} \quad v2 := \begin{bmatrix} -0.5 \\ 0.5 \\ 0 \\ 0 \end{bmatrix} \quad v3 := \begin{bmatrix} 0 \\ 0 \\ 0.5 \\ 0.25 \end{bmatrix} \quad v4 := \begin{bmatrix} 0 \\ 0 \\ -0.5 \\ 0.5 \end{bmatrix}$$

$$y(v) := \begin{bmatrix} \tanh(v_1) \\ \tanh(v_2) \\ \tanh(v_3) \\ \tanh(v_4) \end{bmatrix}$$

$$y1 := y(v1) \quad y2 := y(v2) \quad y3 := y(v3) \quad y4 := y(v4)$$

$$y1 = \begin{bmatrix} 0.462117 \\ 0.244919 \\ 0 \\ 0 \end{bmatrix} \quad y2 = \begin{bmatrix} -0.462117 \\ 0.462117 \\ 0 \\ 0 \end{bmatrix} \quad y3 = \begin{bmatrix} 0 \\ 0 \\ 0.462117 \\ 0.244919 \end{bmatrix} \quad y4 = \begin{bmatrix} 0 \\ 0 \\ -0.462117 \\ 0.462117 \end{bmatrix}$$

Now need to let vx be variable and choose it to give a symmetric W

Define an unknown as vx

$$vx(x1, x2, x3, x4) := \begin{bmatrix} x1 \\ x2 \\ x3 \\ x4 \end{bmatrix} \quad yx(x1, x2, x3, x4) := \begin{bmatrix} \tanh(x1) \\ \tanh(x2) \\ \tanh(x3) \\ \tanh(x4) \end{bmatrix}$$

Inset these vx unknowns into the v vectors so that the 0 assigned entries above will subtract to zero when form vi-vx:

$$vx1(x3, x4) := v1 + \begin{bmatrix} 0 \\ 0 \\ x3 \\ x4 \end{bmatrix} \quad vx2(x3, x4) := v2 + \begin{bmatrix} 0 \\ 0 \\ x3 \\ x4 \end{bmatrix}$$

$$vx3(x1, x2) := v3 + \begin{bmatrix} x1 \\ x2 \\ 0 \\ 0 \end{bmatrix} \quad vx4(x1, x2) := v4 + \begin{bmatrix} x1 \\ x2 \\ 0 \\ 0 \end{bmatrix}$$

There will be two sets of 2x2 matrices to set symmetric with one set being in terms of x1,x2 and the other in terms of x3,x4

Both sets of equations to solve come from

$$W(y1-yx, y2-yx, y3-yx, y4-yx) = G(v1-vx, v2-vx, v3-vx, v4-vx)$$

but this separates into two 2x2 matrices, so separate vx and yx into 2 parts

$$\begin{aligned} vxa(x1, x2) &:= \begin{bmatrix} x1 \\ x2 \end{bmatrix} & vxb(x3, x4) &:= \begin{bmatrix} x3 \\ x4 \end{bmatrix} \\ yxa(x1, x2) &:= \begin{bmatrix} \tanh(x1) \\ \tanh(x2) \end{bmatrix} & yxb(x3, x4) &:= \begin{bmatrix} \tanh(x3) \\ \tanh(x4) \end{bmatrix} \end{aligned}$$

$$V1(x1, x2) := \begin{bmatrix} v1_1 - vxa(x1, x2)_1 & v2_1 - vxa(x1, x2)_1 \\ v1_2 - vxa(x1, x2)_2 & v2_2 - vxa(x1, x2)_2 \end{bmatrix}$$

$$Y1(x1, x2) := \begin{bmatrix} y1_1 - yxa(x1, x2)_1 & y2_1 - yxa(x1, x2)_1 \\ y1_2 - yxa(x1, x2)_2 & y2_2 - yxa(x1, x2)_2 \end{bmatrix}$$

setup for solving for x1 given x2 to make the upper 2x2 submatrix of W symmetric
We have $W=GVY^{-1}$ so form this product and set = its transpose (ignoring the common denominator (= determinant Y1))

$$W12(x1) := G1 \cdot (v2_1 - x1) \cdot (y1_1 - \tanh(x1)) - G1 \cdot (v1_1 - x1) \cdot (y2_1 - \tanh(x1))$$


$$W21(x2) := G2 \cdot (v1_2 - x2) \cdot (y2_2 - \tanh(x2)) - G2 \cdot (v2_2 - x2) \cdot (y1_2 - \tanh(x2))$$

$$f1(x1, x2) := W12(x1) - W21(x2)$$

```

x1min := -1      x1max := 1      x1inc := 0.01
x1 := x1min, x1min + x1inc .. x1max

```



Solve block to determine x_1 given x_2 ; arbitrarily choose x_2 as 0.3

Guess value: $x_2 := 0.3$

$x_1 := 0.5$

Given:

$x_{10} := \text{root}(f1(x1, 0.3), x1)$

$x_{10} = 0.494563$

$x_1 := x_{10}$

Form the upper 2×2 weight matrix which should be symmetric

$$G_{11} := \begin{bmatrix} G_1 & 0 \\ 0 & G_2 \end{bmatrix}$$

$$W_1(x_1, x_2) := G_{11} \cdot V_1(x_1, x_2) \cdot (Y_1(x_1, x_2)^{-1})$$

$$V_1(x_1, x_2) = \begin{bmatrix} 5.436955 \cdot 10^{-3} & -0.994563 \\ -0.05 & 0.2 \end{bmatrix} \quad Y_1(x_1, x_2) = \begin{bmatrix} 4.286609 \cdot 10^{-3} & -0.919948 \\ -0.046394 & 0.170805 \end{bmatrix}$$

$$W_1(x_1, x_2) = \begin{bmatrix} 2.15568 & -0.035206 \\ -0.035213 & 2.1522 \end{bmatrix}$$

Find the input I that goes with these by $I = -Wy + Gv$

$W_1 := W_1(x_1, x_2)$

$$v_{1x} := \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} \quad v_{1x} = \begin{bmatrix} 0.494563 \\ 0.3 \end{bmatrix}$$

$$y_{1x} := \begin{bmatrix} \tanh(x_1) \\ \tanh(x_2) \end{bmatrix} \quad y_{1x} = \begin{bmatrix} 0.457831 \\ 0.291313 \end{bmatrix}$$

$$I_1 := -W_1 \cdot y_{1x} + G_{11} \cdot v_{1x} \quad I_1 = \begin{bmatrix} 0.012446 \\ -0.010841 \end{bmatrix}$$

Next do the same for the lower 2x2 submatrix, now solving for x3,x4

$$V2(x3, x4) := \begin{bmatrix} v3_3 - vx_{\text{b}}(x3, x4)_1 & v4_3 - vx_{\text{b}}(x3, x4)_1 \\ v3_4 - vx_{\text{b}}(x3, x4)_2 & v4_4 - vx_{\text{b}}(x3, x4)_2 \end{bmatrix}$$

$$Y2(x3, x4) := \begin{bmatrix} y3_3 - yx_{\text{b}}(x3, x4)_1 & y4_3 - yx_{\text{b}}(x3, x4)_1 \\ y3_4 - yx_{\text{b}}(x3, x4)_2 & y4_4 - yx_{\text{b}}(x3, x4)_2 \end{bmatrix}$$

setup for solving for x3 given x4 to make the lower 2x2 submatrix of W symmetric


$$W34(x3) := G3 \cdot (v4_3 - x3) \cdot (y3_3 - \tanh(x3)) - G3 \cdot (v3_3 - x3) \cdot (y4_3 - \tanh(x3))$$

$$W43(x4) := G4 \cdot (v3_4 - x4) \cdot (y4_4 - \tanh(x4)) - G4 \cdot (v4_4 - x4) \cdot (y3_4 - \tanh(x4))$$

$$f2(x3, x4) := W34(x3) - W43(x4)$$

$$x3min := -1 \quad x3max := 1 \quad x3inc := 0.01$$

$$x3 := x3min, x3min + x3inc .. x3max$$



Solve block to determine x3 given x4

$$\text{Guess value: } x4 := 0.3$$

$$x3 := 0.5$$

Given:

$$x30 := \text{root}(f2(x3, 0.3), x3)$$

$$x30 = 0.494563 \quad x3 := x30$$

Form the lower 2x2 weight matrix which should be symmetric

$$G22 := \begin{bmatrix} G3 & 0 \\ 0 & G4 \end{bmatrix}$$

$$W2(x3, x4) := G22 \cdot V2(x3, x4) \cdot (Y2(x3, x4)^{-1})$$

$$V2(x3, x4) = \begin{bmatrix} 5.436955 \cdot 10^{-3} & -0.994563 \\ -0.05 & 0.2 \end{bmatrix} \quad Y2(x3, x4) = \begin{bmatrix} 4.286609 \cdot 10^{-3} & -0.919948 \\ -0.046394 & 0.170805 \end{bmatrix}$$

$$W2(x3, x4) = \begin{bmatrix} 2.15568 & -0.035206 \\ -0.035213 & 2.1522 \end{bmatrix}$$

$$W2 := W2(x3, x4)$$

Find the input I that goes with these by $I = -Wy + Gv$

$$\begin{aligned} v3x &:= \begin{bmatrix} x3 \\ x4 \end{bmatrix} & v3x &= \begin{bmatrix} 0.494563 \\ 0.3 \end{bmatrix} \\ y3x &:= \begin{bmatrix} \tanh(x3) \\ \tanh(x4) \end{bmatrix} & y3x &= \begin{bmatrix} 0.457831 \\ 0.291313 \end{bmatrix} \\ I2 &:= -W2 \cdot y3x + G22 \cdot v3x & I2 &= \begin{bmatrix} 0.012446 \\ -0.010841 \end{bmatrix} \end{aligned}$$

Now put all together as the "direct sum" of two 2×2 systems:

$$\begin{aligned} W &:= \begin{bmatrix} W_{1,1,1} & W_{1,1,2} & 0 & 0 \\ W_{1,2,1} & W_{1,2,2} & 0 & 0 \\ 0 & 0 & W_{2,1,1} & W_{2,1,2} \\ 0 & 0 & W_{2,2,1} & W_{2,2,2} \end{bmatrix} & W &= \begin{bmatrix} 2.15568 & -0.035206 & 0 & 0 \\ -0.035213 & 2.1522 & 0 & 0 \\ 0 & 0 & 2.15568 & -0.035206 \\ 0 & 0 & -0.035213 & 2.1522 \end{bmatrix} \\ I &:= \begin{bmatrix} I1_1 \\ I1_2 \\ I2_1 \\ I2_2 \end{bmatrix} & I &= \begin{bmatrix} 0.012446 \\ -0.010841 \\ 0.012446 \\ -0.010841 \end{bmatrix} \end{aligned}$$

Finally check that $Cdv/dt = 0 = g(v) - Wy(v) - Gv + I$ has the desired equilibrium points

First find the actual vectors from given ones plus variable ones

$$veq1 := vx1(x3, x4) \quad veq2 := vx2(x3, x4) \quad veq3 := vx3(x1, x2) \quad veq4 := vx4(x1, x2)$$

$$\begin{aligned} veq1 &= \begin{bmatrix} 0.5 \\ 0.25 \\ 0.494563 \\ 0.3 \end{bmatrix} & veq2 &= \begin{bmatrix} -0.5 \\ 0.5 \\ 0.494563 \\ 0.3 \end{bmatrix} & veq3 &= \begin{bmatrix} 0.494563 \\ 0.3 \\ 0.5 \\ 0.25 \end{bmatrix} & veq4 &= \begin{bmatrix} 0.494563 \\ 0.3 \\ -0.5 \\ 0.5 \end{bmatrix} \\ vx &:= vx(x1, x2, x3, x4) & vx &= \begin{bmatrix} 0.494563 \\ 0.3 \\ 0.494563 \\ 0.3 \end{bmatrix} \end{aligned}$$

Checking that $g(\text{veq}_i)=0$

$$g(v) := W \cdot y(v) - G \cdot v + I$$

$$g(\text{veq1}) = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

$$g(\text{veq2}) = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

$$g(\text{veq3}) = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

$$g(\text{veq4}) = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

$$g(vx) = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

Note that vx has also become an equilibrium point

The neuron outputs for these actual inputs are

$$y(\text{veq1}) = \begin{bmatrix} 0.462117 \\ 0.244919 \\ 0.457831 \\ 0.291313 \end{bmatrix} \quad y(\text{veq2}) = \begin{bmatrix} -0.462117 \\ 0.462117 \\ 0.457831 \\ 0.291313 \end{bmatrix} \quad y(\text{veq3}) = \begin{bmatrix} 0.457831 \\ 0.291313 \\ 0.462117 \\ 0.244919 \end{bmatrix} \quad y(\text{veq4}) = \begin{bmatrix} 0.457831 \\ 0.291313 \\ -0.462117 \\ 0.462117 \end{bmatrix}$$

$$y(vx) = \begin{bmatrix} 0.457831 \\ 0.291313 \\ 0.457831 \\ 0.291313 \end{bmatrix}$$

This is $z(x) = -0.5 \ln((1-z)/(1+z))$

$$z(x) := -0.5 \cdot \ln \left[\frac{(1-x)}{(1+x)} \right] \quad n(x) := \begin{bmatrix} z(x_1) \\ z(x_2) \\ z(x_3) \\ z(x_4) \end{bmatrix}$$

To check $n(y(v))=v$

$$n(y(\text{veq1})) = \begin{bmatrix} 0.5 \\ 0.25 \\ 0.494563 \\ 0.3 \end{bmatrix}$$

$$n(y(\text{veq2})) = \begin{bmatrix} -0.5 \\ 0.5 \\ 0.494563 \\ 0.3 \end{bmatrix}$$

$$n(y(\text{veq3})) = \begin{bmatrix} 0.494563 \\ 0.3 \\ 0.5 \\ 0.25 \end{bmatrix}$$

$$n(y(\text{veq4})) = \begin{bmatrix} 0.494563 \\ 0.3 \\ -0.5 \\ 0.5 \end{bmatrix}$$

$$n(y(vx)) = \begin{bmatrix} 0.494563 \\ 0.3 \\ 0.494563 \\ 0.3 \end{bmatrix}$$

These check our design. Note though that the original vectors $v1, v2, v3, v4$ are not equilibrium points, as seen by the following, but can be obtained by simple projections from the designed equilibria

$$g(v1) = \begin{bmatrix} 0 \\ 0 \\ 0.012446 \\ -0.010841 \end{bmatrix} \quad g(v2) = \begin{bmatrix} 0 \\ 0 \\ 0.012446 \\ -0.010841 \end{bmatrix} \quad g(v3) = \begin{bmatrix} 0.012446 \\ -0.010841 \\ 0 \\ 0 \end{bmatrix} \quad g(v4) = \begin{bmatrix} 0.012446 \\ -0.010841 \\ 0 \\ 0 \end{bmatrix}$$

Determination of the linear transformation to obtain the desired from the result:
Uaes $[v1, v2, v3, v4][veq1, veq2, veq3, veq4]^{-1}$

$$A_{eq} := \begin{bmatrix} veq1_1 & veq2_1 & veq3_1 & veq4_1 \\ veq1_2 & veq2_2 & veq3_2 & veq4_2 \\ veq1_3 & veq2_3 & veq3_3 & veq4_3 \\ veq1_4 & veq2_4 & veq3_4 & veq4_4 \end{bmatrix}$$

$$A_{eq}^{-1} = \begin{bmatrix} -1.555038 & -10.220152 & 2.55674 & 10.226959 \\ -0.858307 & 0.566771 & 0.169637 & 0.678549 \\ 2.55674 & 10.226959 & -1.555038 & -10.220152 \\ 0.169637 & 0.678549 & -0.858307 & 0.566771 \end{bmatrix}$$

$$A := \begin{bmatrix} v1_1 & v2_1 & v3_1 & v4_1 \\ v1_2 & v2_2 & v3_2 & v4_2 \\ v1_3 & v2_3 & v3_3 & v4_3 \\ v1_4 & v2_4 & v3_4 & v4_4 \end{bmatrix}$$

$$A = \begin{bmatrix} 0.5 & -0.5 & 0 & 0 \\ 0.25 & 0.5 & 0 & 0 \\ 0 & 0 & 0.5 & -0.5 \\ 0 & 0 & 0.25 & 0.5 \end{bmatrix}$$

$$Xconvert := A \cdot A_{eq}^{-1}$$

$$Xconvert = \begin{bmatrix} -0.348365 & -5.393461 & 1.193551 & 4.774205 \\ -0.817913 & -2.271652 & 0.724004 & 2.896014 \\ 1.193551 & 4.774205 & -0.348365 & -5.393461 \\ 0.724004 & 2.896014 & -0.817913 & -2.271652 \end{bmatrix}$$

As a check we should get $v1, v2, v3, v4$ back

$$Xconvert \cdot veq1 = \begin{bmatrix} 0.5 \\ 0.25 \\ 0 \\ 0 \end{bmatrix} \quad Xconvert \cdot veq2 = \begin{bmatrix} -0.5 \\ 0.5 \\ 0 \\ 0 \end{bmatrix} \quad Xconvert \cdot veq3 = \begin{bmatrix} 0 \\ 0 \\ 0.5 \\ 0.25 \end{bmatrix}$$

$$Xconvert \cdot veq4 = \begin{bmatrix} 0 \\ 0 \\ -0.5 \\ 0.5 \end{bmatrix}$$

$$Xconvert \cdot vx = \begin{bmatrix} 0.232221 \\ 0.140864 \\ 0.232221 \\ 0.140864 \end{bmatrix}$$